

Correlations between the intensity of fMLP-dependent respiratory burst and cellular fatty acid composition in human neutrophils

P. BELLAVITE, P. GUARINI,* D. BIASI,* A. CARLETTO,* M. T. TREVISAN, P. CARAMASCHI,* L. M. BAMBARA* AND R. CORROCHER* *Institute of Clinical Chemistry and Microscopy and *Medical Pathology, University of Verona, Ospedale Policlinico, Verona, Italy*

Received 6 July 1994; accepted for publication 4 October 1994

Summary. The peripheral blood neutrophils were isolated from a group of normal subjects and their fatty acid composition determined by capillary gas-chromatography. The superoxide (O_2^-) release by the same cell preparations in response to formyl-methionyl-leucyl-phenylalanine was also determined following cytochrome c reduction in a microplate assay. A strong negative correlation was found between C18:2 (linoleic acid) ($r = -0.703$, $P = 0.001$) and C:16:0 (palmitic acid) ($r = -0.569$, $P = 0.009$) and fMLP-stimulated O_2^- release, whereas C20:4 (arachidonic acid) correlated positively ($r = 0.448$, $P = 0.048$). Other fatty acids, namely C12:0, C14:0, C16:1, C18:1, C18:3, C18:4, C20:0, C20:1, C20:2, C20:5, C22:0, C22:1, C22:6, C24:0 and C26:0, were not correlated with O_2^- . No

correlations were found between fatty acid composition and O_2^- release from resting cells and from cells stimulated by phorbol-myristate acetate. These results suggest that the fatty acid composition of blood neutrophils may be a critical factor determining the capability of releasing free radicals in response to formylpeptides. Moreover, since a concomitant increase of arachidonic acid and decrease of its precursor linoleic acid has been found in high-responsive neutrophils, the rate of the enzymes of the arachidonic acid biosynthetic pathway (elongases and desaturases) appear to play an important role.

Keywords: fatty acids, neutrophil priming, superoxide release, signal transduction, phospholipids.

Neutrophils, stimulated by suitable doses of agonists and membrane perturbing agents, are affected by the activation of the enzymatic system NADPH oxidase, which generates excited oxygen species, including superoxide anion (O_2^-) and its derivatives hydrogen peroxide (H_2O_2) and hydroxyl radical ($OH\cdot$). These oxygen species participate in the bactericidal mechanisms of neutrophils, but also in the possible tissue damage caused by the dysregulation of neutrophil activation in a number of pathologies (Sandborg & Smolen, 1988; Bellavite, 1988; Weiss, 1989; Segal & Abo, 1993). Because the duration and the magnitude of the metabolic response are important factors in determining the final balance – positive or unfavourable for the host – of neutrophil activation, these phenomena are subject to a fine and complex regulation at the level of receptors, transduction mechanisms, and NADPH oxidase itself.

In the course of studies on neutrophil activation and priming during inflammation (Bellavite *et al.*, 1992, 1993; Biasi *et al.*, 1993a, b) we noted that the extent of metabolic response to agonist challenge, in terms of superoxide formation, not only varied according to the functional state of the cells (i.e. if they were isolated from blood or from inflammatory exudates), but was quite heterogeneous in a population of normal subjects, even considering the blood cells only. Some of the variability could be due to technical reasons such as the procedures of cell purification, but as a broad distribution of superoxide levels has also been found previously in whole blood measurements (Bellavite *et al.*, 1983), it is more conceivable that the heterogeneity is due to the complex and multiple-step nature of NADPH oxidase activation (McPhail *et al.*, 1984; Maridonneau-Parini *et al.*, 1986; Bellavite, 1988; Bellavite *et al.*, 1990). According to the current knowledge of neutrophil regulation, a conceivable and relevant mechanism explaining interindividual differences of the oxidative metabolism could be the different lipid composition of membranes, due to diet, lifestyle, age, sex,

Correspondence: Professor R. Corrocher, Istituto di Patologia Medica, Ospedale Policlinico, 37134 Verona, Italy.

genetic background, or other unknown factors. In fact, several lines of evidence indicate that lipid composition of the cell membrane may have an important role in the regulation of the respiratory burst of neutrophils. Certain saturated fatty acids (myristate in particular) and *cis*-unsaturated fatty acids (arachidonate in particular) elicit production of O_2^- in intact cells (Kakinuma, 1974; Curnutte *et al*, 1984). A direct role of lipids in the regulation of activation or activity of the free-radicals-generating oxidase has been suggested by the stimulating effect on the solubilized and isolated enzyme exhibited by phosphatidylethanolamine (Gabig & Babior, 1979), by phosphatidylserine (Tamura *et al*, 1988) and by phosphatidic acid (Bellavite *et al*, 1988). Moreover, the NADPH oxidase in cell-free systems is activated by arachidonic acid and other unsaturated fatty acids (Bromberg & Pick, 1984). Finally, it has been shown recently that arachidonic acid acts also directly on the assembled NADPH oxidase in the membranes and in the endosomes and increases its catalytic activity by increasing the affinity for NADPH (Rubinck & Levy, 1993).

These data and observations prompted us to investigate the relationship between lipid composition and oxidative metabolic responses in a simple and straightforward way, i.e. by measuring, in a group of normal human subjects, the amount of neutrophil O_2^- production and the amount of the most representative species of fatty acids contained in the same cells. To the best of our knowledge, no similar data are available in the literature.

Neutrophils were isolated from peripheral blood, then divided in two aliquots, one of which was used for the determination of fatty acids, the other was used for superoxide production in a microplate assay. Superoxide production was measured on resting cells, on cells stimulated by an agonist that acts through specific membrane receptors, namely *n*-formyl-methionyl-leucyl-phenylalanine (FMLP) and on cells stimulated by phorbol-myristate acetate (PMA), an agent which is known to perform its effects through direct stimulation of protein kinase C. Both positive and negative correlations between the two considered parameters emerged from our study.

METHODS

Reagents. FMLP, PMA and reference fatty acids were purchased from Sigma Chemical Company, St Louis, Mo.; cytochrome *c* from Boehringer, Mannheim, Germany; purified serum bovine albumin and human albumin from Behring Institut, Marburg, Germany; Percoll was from Pharmacia, Uppsala, Sweden. Sterile 96-well microtitre plates with flat-bottomed wells (Linbro type) were from Flow Laboratories. The microplates were pre-coated with fetal bovine serum (Flow Laboratories) in order to abolish nonspecific cell activation as described (Bellavite *et al*, 1992). Hanks balanced salt solution (without calcium and magnesium) (HBSS) and Dulbecco's phosphate-buffered saline (PBS) were from Gibco Ltd, Paisley, Scotland; the composition of HBSS was 0.4 g/l KCl, 0.06 g/l KH_2PO_4 , 8 g/l NaCl, 0.09 g/l Na_2HPO_4 , 1 g/l D-glucose (pH 7.4); the composition of PBS was 0.2 g/l KCl, 0.2 g/l KH_2PO_4 ,

0.047 g/l $MgCl_2$, 8 g/l NaCl, 1.15 g/l Na_2HPO_4 (pH 7.4). Other materials and reagents were of the highest purity available.

Cell preparation. Neutrophils were obtained from blood of normal human volunteers by centrifugation over discontinuous Percoll gradients (Metcalfe *et al*, 1986). Ethylene diamine tetraacetate-anticoagulated blood samples were centrifuged over Percoll gradients for 20 min at 1800 rpm with a Sorvall T6000B centrifuge. Neutrophils were recovered as a broad band at the 73%–62% interface, were diluted with 1 volume of PBS, and centrifuged for 10 min at 1200 rpm. The pellet was usually slightly contaminated by erythrocytes, that were lysed by a brief hypotonic shock: cells were suspended in 5 ml of 0.2% NaCl for 20 s, then the isotonicity was restored by addition of 5 ml of 1.6% NaCl and right pH was restored by addition of 2 ml of PBS. After hypotonic lysis of contaminating erythrocytes and two washings with PBS, the cells (>95% neutrophils, >99% viable as judged by trypan blue exclusion test) were finally suspended in HBSS, containing 0.2% human serum albumin (H-A), and kept at room temperature until use. A few minutes before use, 100× concentrated solutions of $CaCl_2$ and $MgSO_4$ were added to the cell suspensions at the final concentration of 0.5 mM and 1 mM respectively.

Superoxide measurement. A microplate assay of O_2^- production was performed according to previously reported procedures (Bellavite *et al*, 1992, 1993), with the following modifications. The assay medium was H-A supplemented with 0.5 mM $CaCl_2$ and 1 mM $MgSO_4$ (H-ACM). The microplate wells were supplemented with 25 μ l of 0.6 mM cytochrome *c* and with either 25 μ l of the stimulant dissolved in H-ACM (FMLP or PMA at the final concentration of 5×10^{-7} M and 10 ng/ml respectively) or with 25 μ l of H-ACM (resting assays). Assays were currently done in triplicate for each experimental condition. The plate was brought to 37°C, and 50 μ l of the neutrophil suspension (2×10^5 cells), pre-warmed at 37°C, were added to each well. The plate was then incubated for 10 min in a humidified thermostat at 37°C. When required, the plate was rapidly transferred into a microplate reader (Reader 400, SLT Labs Instruments) and the reduction of cytochrome *c* was measured at 550 nm using 540 nm as reference wavelength, and using 0.037 optical density units as standard for 1 nmol of reduced cytochrome *c* (Bellavite *et al*, 1992).

Fatty acid determination. Analysis of fatty acids was performed on aliquots of neutrophil preparations containing 10^7 cells, by extraction of total lipids with 4.5 ml of isopropanol/chloroform (11/7, v/v) added with 0.45 mM 2,6-di-*ter*-*p*-cresol (BHT) as antioxidant. A gas chromatographic (Hewlett Packard 5890 chromatograph; Hewlett Packard, Palo Alto, Calif., U.S.A.) method was employed as previously described (Girelli *et al*, 1992), based on fatty acid direct transesterification technique (Lepage & Roy, 1986). Analysis were performed in duplicate on each sample. Peak identification and quantification were made with reference fatty acids commercially available. As internal standard, heptadecanoic acid (C17:0) was used. The heights of the peaks were measured and the quantification was performed

with the aid of a PC Vectra (QS/16S equipped with the software HP-3365 ChemStation Hewlett Packard) working in Microsoft Windows 3.0. Fatty acid composition data were expressed as mg/100 mg fatty acid methyl esters. The mean percentage recovery was 90.6 ± 2.7 (range 85–94%).

Data analysis. Statistical analysis was performed on a PC Apple Macintosh SE/30 equipped with the software Systat 5.0, working in Windows 3.0 (Microsoft). Values are expressed as mean \pm standard deviation and the correlation between different parameters was studied using the Pearson correlation coefficient.

RESULTS

The mean fatty acid compositions of neutrophils from the group of 20 normal volunteers used in this study are given in Table I. It can be seen that 44% of fatty acids were saturated, the most represented being palmitic and stearic; 30% were monounsaturated, represented almost exclusively by oleic acid; a quarter of fatty acids were polyunsaturated and, among these, 9% was represented by linoleic acid and 13% by arachidonic acid.

The O_2^- release by the neutrophils (nmol/10 min/ 10^6 cells, $n = 20$) was as follows: resting cells, 0.92 ± 0.82 (range 0.0–3.0); fMLP-activated cells, 11.04 ± 3.55 (range 6.0–17.0); PMA-activated cells 28.33 ± 9.4 (range 19.5–49.6). The fatty acid content of blood neutrophils was then plotted against the production of superoxide. The oxidative metabolism of resting (unstimulated) cells was so low that no possible correlations could be calculated with the fatty acids content. Therefore the only consistent data were obtained on fMLP- and PMA-activated cells and the correlation drawn was between fatty acid content and the activation of the metabolism. Among the fatty acids species reported in Table I, the majority were not correlated with oxidative metabolism and nor was correlation found with the total content of unsaturated and saturated fatty acids (not shown). However, three fatty acid species showed a significant correlation with fMLP-activated respiratory burst

Table I. Fatty acid content of neutrophils from 20 normal subjects.

	Total fatty acids (w/w) (%) \pm SD
Saturated	
Total	44.40 ± 2.35
C12:0	0.27 ± 0.15
C14:0	0.77 ± 0.23
C16:0 (palmitic)	21.35 ± 1.19
C18:0 (stearic)	18.84 ± 2.19
C20:0	0.38 ± 0.02
C22:0	1.28 ± 0.19
C24:0	1.92 ± 0.26
C26:0	0.37 ± 0.14
Monounsaturated	
Total	29.91 ± 2.74
C16:1w7	0.52 ± 0.24
C18:1w9 (oleic)	27.94 ± 2.19
C20:1w9	1.01 ± 0.23
C22:1w9	1.34 ± 0.51
Polyunsaturated	
Total	24.93 ± 1.93
C18:2w6 (linoleic)	9.32 ± 1.71
C18:4w3	1.22 ± 0.24
C20:2w6	0.98 ± 0.29
C20:4w6 (arachidonic)	12.81 ± 1.63
C20:5w3	0.60 ± 0.03
C22:6w6	1.26 ± 0.32

(Fig 1). These were palmitic acid and linoleic acid which showed a negative correlation, and arachidonic acid which showed a weak, but significant, positive correlation. In other words, the superoxide production was higher in cells having a low content of palmitic and linoleic acid and a high content of arachidonic acid.

It should be pointed out that the determinations of superoxide production and of fatty acids were done by two




Fig 1. Linear regression plot of fatty acid content and fMLP-stimulated O_2^- release in human neutrophils.




Fig 2. Correlation between arachidonic acid and linoleic acid content in human neutrophils.

independent teams who were unaware of the results obtained by the other. The correlations were calculated at the end, when all the experimental data had been collected.

When the same calculation was done plotting the fatty acid content against the superoxide produced in response to PMA, no correlation was found with any of the fatty acids determined. The data concerning the major fatty acids are: C16:0, $r: -0.188$, $P: 0.287$; C18:2, $r: -0.038$, $P: 0.830$; C20:4, $r: 0.335$, $P: 0.052$.

Since linoleic acid is a precursor of arachidonic acid, it was of interest to see if the cellular content of the two fatty acids were inversely correlated. Fig 2 shows that this is the case: arachidonic acid showed a net tendency to be higher when the linoleic acid was lower, as expected.

DISCUSSION

Neutrophils can be present in various activity states, varying from totally resting, to primed (pre-conditioned to enhanced responses), to fully activated. The neutrophils may also be present in de-activated, or adapted, or desensitized states, which are low-response states, induced by various mechanisms which may down-regulate either receptors or the effector systems. The various activity states of these cells are relevant for the non-specific immunological surveillance against microbial invaders and also for the potentially harmful role played by these cells in pathological processes such as rheumatoid arthritis, adult respiratory distress syndrome, post-ischaemic tissue injury, immune complex diseases, etc. Many compounds such as cytokines, bacterial products, opioid peptides, hormones capable of regulating the neutrophil responses and, in particular, the extent of the free-radical generating respiratory burst have been identified, but less is known regarding the mechanisms which are

involved in the changes of sensitivity or of responsiveness at cellular level.

In this paper we have found that the production of superoxide anion in response to fMLP is in direct correlation with the arachidonate content of neutrophils and in inverse relationship with the linoleate and palmitate content, suggesting that the lipid composition of the cells should be regarded as a potential regulatory factor involved in the determination of cell activity. It is well known that phospholipids stimulate a variety of enzyme-catalysed oxidative reactions, including the NADPH oxidase (Curnutte *et al.* 1984; Gabig & Babior, 1979; Tamura *et al.* 1988). Changes in the phospholipid and fatty acid composition of plasma membranes under some conditions could modulate the function of the activated oxidase, thereby exerting regulatory effects on superoxide generation.

Finding a correlation does not establish a causal relationship, and further studies are necessary in order to provide an explanation for the enhanced responses in the cells which are more rich in arachidonic acid and less rich in linoleic acid. A possible simple explanation of our findings could be the increased availability of arachidonate, which is well known to be a NADPH oxidase activator, possibly through a charge-dependent process (Bromberg & Pick, 1984; Miyahara *et al.* 1987; Bellavite *et al.* 1988; McPhail *et al.* 1993). The effects of arachidonic acid seem to be quite specific, because two other unsaturated fatty acids, linoleic and oleic acid, and a saturated one, stearic acid, are much less potent (Rubinek & Levy, 1993). Arachidonic acid is therefore considered the immediate signal for the activation of the oxidase (Henderson *et al.* 1993).

Our data do not exclude that the lipid composition could also exert a regulating function on the fMLP receptor and signal transduction, rather than a direct effect on the terminal NADPH oxidase. This possibility seems to be consistent with the fact that the fatty acid composition affects the responses to fMLP and not those to PMA, an agent which bypasses membrane receptors and many steps of the transduction cascade. We have previously observed that neutrophil priming during an inflammatory process is factor-specific and does not affect the response to PMA (Biasi *et al.* 1993a). Others have reported that the inhibition of phospholipase A2 results in the inhibition of O_2^- generation induced by opsonized zymosan, but not by PMA, suggesting that in the latter case the arachidonic acid is not involved (Sandborg & Smolen, 1988).

In addition to a role in the direct activation of the oxidase, arachidonic acid has been suggested as playing a role in the priming phenomenon (DiPersio *et al.* 1988; Bauldry *et al.* 1991; Corey & Rosoff, 1991; Forehand *et al.* 1993) and this may be the case in our model system, where the cells which have higher arachidonic acid content do not produce superoxide in the resting state, but are in some way primed for an enhanced response to fMLP. Others have shown that incorporation of unsaturated fatty acids into phospholipids of macrophages leads to an increase of O_2^- production and to an increase of phospholipase A2 activity (Flesch & Ferber, 1986).

Linoleic acid and palmitic acid showed a strong negative

correlation with fMLP-stimulated superoxide production. This evidence is difficult to explain, because no studies on the regulation of the oxidase by these fatty acids are available. An effect based on changes of membrane fluidity seems unlikely, because the total content of unsaturated and saturated fatty acids did not correlate with the respiratory burst. At least for the linoleate, it is tempting to speculate that the low content of this fatty acid moiety in high-responder cells is directly related to the increase of the metabolic pathway leading to the formation of arachidonate, of which the linoleate is the main precursor.

In the polyunsaturated fatty acid biosynthetic pathways, arachidonic acid is generated starting from linoleic acid, which undergoes two desaturation steps and one elongation step. Seeing that our data show that arachidonic acid and linoleic acid in our subject population are inversely correlated, the extent of transformation of linoleic in arachidonic could be an important factor regulating the intensity of the response to fMLP of neutrophils. If this case, the activity of desaturase enzymes and of elongase may have an important role in the overall regulation of neutrophil activation.

Whatever the mechanism involved behind the correlations reported here may be, our data confirm and reinforce the lines of study pointing to the possibility of manipulation of neutrophil functions by changes in fatty acid composition. Membrane lipid modifications can be achieved *in vitro* by controlling the lipid environment of the cell (Lou, 1984; Fleisch & Ferber, 1986) and *in vivo* by dietary means. For instance, (a) macrophage arachidonate metabolism in mice is decreased by a fish oil diet and this diet is beneficial because it retards the development of experimental amyloidosis (Cathcart *et al.*, 1987), (b) alimentation with a diet rich in fish oil reduced the superoxide response to fMLP in guinea-pig peritoneal leucocytes compared with a diet rich in oleic acid (Fletcher & Ziboh, 1990), and (c) diets enriched with soy phosphatidylcholine can increase the arachidonate concentration in the membranes, the arachidonate release and the phagocytosis and killing of *Candida albicans* (Jannace *et al.*, 1992). In the latter case, it has been suggested that the increase of arachidonate content obtained by dietary supplementation may have a potential immunostimulatory role in humans who are at risk of infections.

ACKNOWLEDGMENTS

This work was supported by a grant from Ministero Università Ricerca Scientifica e Tecnologica and from Public Health Department, Regione Veneto. The valuable advice of Vittorina Della Bianca and the skilful technical assistance of Fabio Poli are also acknowledged.

REFERENCES

- Bauldry, S.A., McCall, C.E., Cousart, S.L. & Bass, D.A. (1991) Tumor necrosis factor- α priming of phospholipase A2 activation in human neutrophils: an alternative mechanism of priming. *Journal of Immunology*, **146**, 1277–1285.
- Bellavite, P. (1988) The superoxide-forming enzymatic system of phagocytes. *Free Radical Biology and Medicine*, **4**, 225–261.
- Bellavite, P., Chirumbolo, S., Lippi, G., Guzzo, P. & Santonastaso, C. (1993) Homologous priming in chemotactic peptide stimulate neutrophils. *Cell Biochemistry and Function*, **11**, 93–100.
- Bellavite, P., Chirumbolo, S., Mansoldo, C., Andini, G. & Dri, P. (1992) A simultaneous assay for oxidative metabolism and adhesion of human neutrophils. Evidence for correlations and dissociations of the two responses. *Journal of Leukocyte Biology*, **51**, 329–335.
- Bellavite, P., Corso, F., Dusi, S., Grzeskowiak, M., Della Bianca, V. & Rossi, F. (1988) Activation of a NADPH-dependent superoxide production in plasma-membrane extracts of pig neutrophils by phosphatidic acid. *Journal of Biological Chemistry*, **263**, 8210–8214.
- Bellavite, P., Dri, P., Della Bianca, V. & Serra, M.C. (1983) The measurement of superoxide anion production by granulocytes in whole blood: a clinical test for the evaluation of phagocyte function and serum opsonic capacity. *European Journal of Clinical Investigation*, **13**, 363–368.
- Bellavite, P., Serra, M.C., Bazzoni, F., Miron, S. & Dusi, S. (1990) Triggering and regulation of the free radical production by phagocytes. *Free Radicals, Lipoproteins and Membrane Lipids* (ed. by A. Crastes de Paulet, L. Douste Blazy and R. Paoletti), pp. 23–44. Plenum Press, New York.
- Biasi, D., Bambara, L.M., Carletto, A., Caraffi, M., Serra, M.C., Chirumbolo, S. & Bellavite, P. (1993a) Factor-specific changes of the oxidative metabolism of exudate human neutrophils. *Inflammation*, **17**, 13–23.
- Biasi, D., Bambara, M.L., Carletto, A., Caramaschi, P., Serra, M.C., Guzzo, P., Santonastaso, C. & Bellavite, P. (1993b) Molecular mechanisms of priming: exudate human neutrophils are primed to Met-Leu-Phe but have normal intracellular calcium and cAMP responses. *Bulletin of Molecular Biology Medicine*, **18**, 225–235.
- Bromberg, Y. & Pick, E. (1984) Unsaturated fatty acids stimulate NADPH-dependent superoxide production by cell-free system derived from macrophages. *Cellular Immunology*, **88**, 213–221.
- Cathcart, E.S., Leslie, C.A., Meydani, S.N. & Hayes, K.C. (1987) A fish oil diet retards experimental amyloidosis, modulates lymphocyte function, and decreases macrophage arachidonate metabolism in mice. *Journal of Immunology*, **139**, 1850–1854.
- Corey, S.J. & Rosoff, P.M. (1991) Unsaturated fatty acids and lipoxygenase products regulate phagocytic NADPH oxidase activity by a nondetergent mechanism. *Journal of Laboratory and Clinical Medicine*, **118**, 343–351.
- Curnutte, J.T., Badwey, J.A., Robinson, J.M., Karnovsky, M.J. & Karnovsky, M.L. (1984) Studies on the mechanism of superoxide release from human neutrophils stimulated with arachidonate. *Journal of Biological Chemistry*, **259**, 11851–11857.
- DiPersio, J.F., Billing, P., Williams, R. & Gasson, J.C. (1988) Human granulocyte-macrophage colony-stimulating factor and other cytokines prime human neutrophils for enhanced arachidonic acid release and leukotriene B4 synthesis. *Journal of Immunology*, **140**, 4315–4322.
- Fleisch, I. & Ferber, E. (1986) Effect of cellular fatty acid composition on the phospholipase A2 activity of bone marrow-derived macrophages, and their ability to induce lucigenin-dependent chemiluminescence. *Biochimica et Biophysica Acta*, **889**, 6–14.
- Fletcher, M.P. & Ziboh, V.A. (1990) Effects of dietary supplementation with eicosapentaenoic acid or gamma-linoleic acid on neutrophil phospholipid fatty acid composition and activation responses. *Inflammation*, **14**, 585–597.
- Forehand, J.R., Johnston, R.B., Jr & Bomalaski, J.S. (1993) Phospholipase A2 activity in human neutrophils: stimulation by

- lipopolysaccharide and possible involvement in priming for an enhanced respiratory burst. *Journal of Immunology*, **151**, 4918–4925.
- Gabig, T.G. & Babior, B.M. (1979) the O_2^- -forming oxidase responsible for the respiratory burst in human neutrophils: properties of the solubilized enzyme. *Journal of Biological Chemistry*, **254**, 9070–9074.
- Girelli, D., Azzini, M., Olivieri, O., Guarini, P., Trevisan, M.T., Lupo, A., Bernich, P., Panzetta, G. & Corrocher, R. (1992) Red blood cells and platelet membrane fatty acids in non-dialyzed and dialyzed uremics. *Clinica Chimica Acta*, **211**, 155–166.
- Henderson, L.M., Moule, S.K. & Chappel, J.B. (1993) The immediate activator of the NADPH oxidase is arachidonate, not phosphorylation. *European Journal of Biochemistry*, **211**, 157–162.
- Jannace, P.W., Lerman, R.H., Santos, J.I. & Vitale, J.J. (1992) Effects of oral soy phosphatidylcholine on phagocytosis, arachidonate concentrations, and killing by human polymorphonuclear leukocytes. *American Journal of Clinical Nutrition*, **56**, 599–603.
- Kakinuma, K. (1974) Effects of fatty acids on the oxidative metabolism of leukocytes. *Biochimica et Biophysica Acta*, **348**, 76–85.
- Lepage, G. & Roy, C.C. (1986) Direct transesterification of all classes of lipids in a one-step reaction. *Journal of Lipid Research*, **27**, 114–120.
- Lou, J.S. (1984) Incorporation of arachidonic acid and eicosapentaenoic acid into phospholipids by polymorphonuclear leukocytes in vitro. *Lipids*, **19**, 573–577.
- Maridonneau-Parini, I., Tringale, S.M. & Tauber, A.I. (1986) Identification of distinct activation pathways of the human neutrophil NADPH-oxidase. *Journal of Immunology*, **137**, 2925–2929.
- McPhail, L.C., Clayton, C.C. & Snyderman, R. (1984) The NADPH oxidase of human polymorphonuclear leukocytes: evidence for regulation by multiple signals. *Journal of Biological Chemistry*, **259**, 5768–5775.
- McPhail, L.C., Qualliotine-Mann, D., Agwu, D.E. & McCall, C.E. (1993) Phospholipases and the activation of the NADPH oxidase. *European Journal of Haematology*, **51**, 294–300.
- Metcalf, J.A., Gallin, J.I., Nauseef, W.M. & Root, R.K. (1986) *Laboratory Manual of Neutrophil Function*. Raven Press, New York.
- Miyahara, M., Watanabe, N., Okimasu, E. & Utsumi, K. (1987) Charge-dependent regulation of NADPH oxidase activity in guinea-pig polymorphonuclear leukocytes. *Biochimica et Biophysica Acta*, **929**, 253–262.
- Rubinek, T. & Levy, R. (1993) Arachidonic acid increases the activity of the assembled NADPH oxidase in cytoplasmic membranes and endosomes. *Biochimica et Biophysica Acta*, **1176**, 51–58.
- Sandborg, R.R. & Smolen, J.E. (1988) Biology of disease: early biochemical events in leukocyte activation. *Laboratory Investigation*, **59**, 300–320.
- Segal, A.W. & Abo, A. (1993) The biochemical basis of the NADPH oxidase of phagocytes. *Trends in Biochemical Sciences*, **18**, 43–47.
- Tamura, M., Tamura, T., Tyagi, S.R. & Lambeth, J.D. (1988) The superoxide-generating respiratory burst oxidase of human neutrophil plasma membrane. Phosphatidylserine as an effector of the activated enzyme. *Journal of Biological Chemistry*, **263**, 17621–17626.
- Weiss, S.J. (1989) Tissue destruction by neutrophils. *New England Journal of Medicine*, **320**, 365–376.