



Replications of fundamental research models in ultra high dilutions 1994 and 2015 – update on a bibliometric study

Peter Christian Endler^{1,*}, Paolo Bellavite², Leoni Bonamin³, Tim Jäger^{4,5,6} and Sintia Mazon¹

¹Interuniversity College for Health and Development Graz/Castle of Seggau, Austria

²University of Verona, Italy

³Universidade Paulista, Sao Paulo, Brazil

⁴University of Witten/Herdecke, Germany

⁵Institute of Complementary Medicine IKOM, University of Bern, Switzerland

⁶University of Freiburg, Germany

Introduction: This paper focuses exclusively on experimental models with ultra high dilutions (i.e. beyond 10^{-23}) that have been submitted to replication scrutiny. It updates previous surveys, considers suggestions made by the research community and compares the state of replication in 1994 with that in 2015.

Methods: Following literature research, biochemical, immunological, botanical, cell biological and zoological studies on ultra high dilutions (potencies) were included. Reports were grouped into *initial studies*, *laboratory-internal*, *multicentre* and *external* replications. Repetition could yield either *comparable*, or *zero*, or *opposite* results. The null-hypothesis was that test and control groups would not be distinguishable (zero effect).

Results: A total of 126 studies were found. From these, 28 were initial studies. When all 98 replicative studies were considered, 70.4% (i.e. 69) reported a result comparable to that of the initial study, 20.4% (20) zero effect and 9.2% (9) an opposite result. Both for the studies until 1994 and the studies 1995–2015 the null-hypothesis (dominance of zero results) should be rejected. Furthermore, the odds of finding a *comparable* result are generally higher than of finding an *opposite* result. Although this is true for all three types of replication studies, the fraction of *comparable* studies diminishes from laboratory-internal (total 82.9%) to multicentre (total 75%) to external (total 48.3%), while the fraction of *opposite* results was 4.9%, 10.7% and 13.8%. Furthermore, it became obvious that the probability of an external replication producing comparable results is bigger for models that had already been further scrutinized by the initial researchers.

Conclusions: We found 28 experimental models which underwent replication. In total, 24 models were replicated with comparable results, 12 models with zero effect, and 6 models with opposite results. Five models were *externally* reproduced with comparable results. We encourage further replications of studies in order to learn more about the model systems used. *Homeopathy* (2015) 104, 234–245.

Keywords: Review; Basic research; Homeopathy; Ultra high dilution; Replication

Introduction

There are several thousand references on fundamental research in homeopathy, including hundreds of references on extreme dilutions. This paper focuses exclusively on experimental models with ultra high dilutions (i.e. beyond 10^{-23}) that have been submitted to replication scrutiny.^{1–101} It follows on from a previous survey of

*Correspondence: Peter Christian Endler, Interuniversity College for Health and Development Graz/Castle of Seggau, Petrifelderstr. 4, A-8042 Graz, Austria.

E-mail: college@inter-uni.net

Received 24 June 2015; revised 2 October 2015; accepted 13 October 2015

2009¹⁰² which gave an overview of fundamental biochemical and biological studies that used high homeopathic potencies and that had been subjected to laboratory-internal, multicentre or external replication trials. Physicochemical or clinical studies were not included, nor studies on dilutions below 10^{-23} , nor studies in relation to which no attempt of replication could be found in literature.

The studies under survey were grouped into broadly defined clusters according to the methodology employed (see below [Methods](#)).

Apart from being a mere update this paper considers suggestions made by the research community in response to the first publication¹⁰² regarding the literature surveyed as well as its clustering, evaluation and discussion. It also compares the state of replication in 1994, when the anthology ‘Ultra High Dilution’ was published,⁹⁹ with that in 2015. Furthermore, with regard to the models presented, we tried to determine whether it makes sense to pursue laboratory-internal and multicentre replication research as a means of mitigating the probability of external replication studies producing zero results.

Methods

Literature search

Sources of information were reviews,^{99,102–113} personal contact with members of the homeopathic research community, and the MEDLINE (www.PubMed.gov) and HOMBREX (www.carstens-stiftung.de) databases. Allowed literature sources were publications (in peer-reviewed and not peer-reviewed journals, book sections and books) as well as unpublished academic papers. As a rule, unpublished papers were disregarded wherever published papers on the same study were available. Especially from 2010 to 2015, we focused on PubMed listed publications. Although we have done what seemed possible to identify all relevant studies, the annotated bibliography presented here does not claim to be exhaustive.

Inclusion criteria

We included biochemical, immunological, botanical, cell biological and zoological studies on ultra high, homeopathically prepared dilutions (potencies), i.e. $\geq 24x$ ($=10^{-24}$) or 12c ($=100^{-12}$). Studies published after 1940 were required to report evaluation of results by statistical methods (minimum requirement: mean or median, number n of samples, standard deviation or standard error, OR, number N of samples, level of significance of a statistical test).

To be included the experiment had to have been repeated. Replications were formally considered as such whenever it was possible to find

- at least two publications by the same initial working group, including a follow-up trial of an initial publication (laboratory-internal replication) or
- at least one publication reporting on a multicentre trial (independent experiments in different locations/labora-

tories, organized by one study coordinator, normally from the initial team), or

- at least one publication by the initial workgroup and one with external authorship, both dealing with the same experimental model.

Furthermore, replication was considered as such when a later study dealt with the same biological system and the same potentized substance as an earlier one. Within such clusters, however, a certain degree of deviation was accepted with regard to the biological system (e.g. the use of *Chlorella vulgaris* or *Chlorella pyrenoidosa*), the potency level (e.g. 25x or 30x) and potency type (decimal (x) or centesimal (c)) and the nature of the control (e.g. prepared step by step or not, succeeded or not, or type not mentioned).

One and the same publication could refer to the results of more than one study. Where numbers of studies are quoted in connection with multicentre trials they refer to the number of trials in different locations/laboratories. Among the initial studies, one of the researchers involved was always considered as the ‘initial’ researcher. When their name could not be identified from the publication, the first author’s name was mentioned. Apart from the main publications, four publications giving additional information were cited.^{98–101}

Thus, we extracted all studies from the included publications and grouped them into experimental models.

Studies (i.e. initial and repeated studies) were further grouped according to results achieved:

- Initial studies: as an inclusion criterion to be candidates for replication trials, these had shown a significant difference between test and control group, e.g. enhancing growth
- Repeated studies, the results of which were *consistent* with the initial study, i.e. where a *comparable* result (in the same direction, e.g. enhancing growth) was found
- Repeated studies, where *no* difference between test and control group was found (zero effect)
- Repeated studies, the results of which were *opposite* to the initial study, i.e. when results were *different* in direction (e.g. decreasing instead of increasing).

The null-hypothesis was that test and control groups would not be distinguishable, i.e. there would be no result of treatment with the potency (zero effect). In this survey paper, we focused on a graphical representation of the data rather than statistical calculations. Raw data for further analysis are given in [Table 1](#) (see [Results](#)).

Results

A total of 126 studies were found. [Figure 1](#) shows the proportion of models and studies.

Numbers of studies performed until 1994¹⁰² and in the period from 1995 to 2015 were: on enzymes 2 + 7, on cultured cells 0 + 2, on plants 4 + 30, on immune cells 12 + 16, on isolated organs 0 + 4, on amphibians/fish 8 + 18 and on rats/mice 9 + 14.

Table 1 Fundamental models and studies on ultra high dilutions. Studies were classified according to the type of replication (initial, laboratory-internal, multicentre, external) and to the results achieved (comparable, zero, opposite)

<i>biochemistry: enzyme diastase & mercury chloride</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Persson 1932 ¹			
<i>multicentre</i>				
<i>external</i>			Bluth 2005 ²	Boyd 1954 ³
<i>resumee: increase of enzyme reaction in 1 study, decrease in 1, zero result in 1</i>				
<i>biochemistry: enzyme acid phosphatase & ubiquinone</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Harisch 1997 ⁴	Harisch 1999 ⁵		
<i>multicentre</i>				
<i>external</i>				
<i>resumee: decrease of enzyme reaction in both studies</i>				
<i>biochemistry: enzyme acid phosphatase & cAMP</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Harisch 1998 ⁶	Harisch 1999 ⁷		
<i>multicentre</i>				
<i>external</i>				
<i>resumee: decrease of enzyme reaction in both studies</i>				
<i>biochemistry: enzyme alpha amylase & mercury chloride</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Sukul 2002 ⁸			
<i>multicentre</i>				
<i>external</i>			Bluth 2005 ²	
<i>resumee: increase of enzyme reaction in the initial study, zero result in the repetition</i>				
<i>cultured mammalian cells: neuroblastoma cells & tumour necrosis factor alpha</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Carmine 1997 ⁹			
<i>multicentre</i>				
<i>external</i>			Herberth 1999 ¹⁰	
<i>resumee: increase of H2O2 production in the initial study, zero result in the repetition</i>				
<i>plants: algae Chlorella & copper sulphate</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Graviou 1971 ¹¹			
<i>multicentre</i>				
<i>external</i>			Moss 1977 ¹²	
<i>resumee: growth stimulation of poisoned algae in the initial study, zero result in the repetition</i>				
<i>plants: wheat seedlings & silver nitrate</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Kolisko 1926 ¹³			
<i>multicentre</i>				
<i>external</i>		Pongratz 1994 ¹⁴	Endler 1998 ¹⁵	
		Pongratz 1998 ¹⁵		
		Nogrask 1998 ¹⁵		
<i>resumee: increase of stalk growth in 4 studies, zero result in 1 study</i>				
<i>plants: arsenic poisoned wheat seedlings & arsenicum album</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Betti 1997 ¹⁶	Brizzi 2000 ¹⁷		
		Brizzi 2005 ¹⁸		
		Nani 2007 ¹⁹		
		Brizzi 2011 ²⁰		
<i>multicentre</i>				
<i>external</i>			Lahnstein 2009 ²¹	Binder 2005 ²²
				Lahnstein 2009 ²¹
<i>resumee: stimulation of growth and germination rate, decrease of variance in 5 studies, decrease of growth and germination rate in 2 studies, increase of variance in 1 study</i>				
<i>plants: wheat seedlings (stalk growth) & gibberellic acid</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Pfleger 2008 ²³	Hofäcker 2008 ²³		
		Reich 2009 ²⁴		
		Hribar 2013 ²⁵		
<i>multicentre</i>				Reischl 2009 ²⁴
<i>external</i>				Thieves 2009 ²⁴
<i>resumee: decrease of stalk growth in 4 studies, increase in 2</i>				
<i>hypothesis: decrease in growth season, increase in winter</i>				
<i>plants: wheat seedlings (germination) & gibberellic acid</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Schiestl 2010 ²⁶			
<i>multicentre</i>		Hartung 2010 ²⁶	Matzer 2012 ²⁷	Hofstätter 2012 ²⁷
		Schwärzler 2012 ²⁷	Seunig 2012 ²⁷	
<i>external</i>				
<i>resumee: decrease of germination in 3 studies, increase in 1 study, zero result in 1 study</i>				
<i>plants: duckweed & gibberellic acid</i>				

Table 1. (Continued)

	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Scherr 2009 ²⁸			Majewsky 2014 ²⁹
<i>multicentre</i>				
<i>external</i>				
<i>resumee: increase of growth in 1 study, decrease in 1</i>				
<i>plants: cress growth and biocrystallisation & stannum metallicum</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Andersen 2012 ³⁰	Andersen 2015 ³¹		
<i>multicentre</i>		Doesburg 2012 ³⁰		
		Doesburg 2015 ³¹		
<i>external</i>				
<i>resumee: additive-specific crystallisation patterns in all 4 studies</i>				
<i>isolated immune cells: basophils & antiserum against IgE</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Davenas 1988 ³²	Benveniste 1991 ³³		
<i>multicentre</i>				
<i>external</i>			Ovelgönne 1992 ³⁴	
			Hirst 1993 ³⁵	
<i>resumee: reduction of degranulation in 2 studies, zero result in 2</i>				
<i>isolated immune cells: basophils & apis mellifica</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Poitevin 1986 ³⁶	Poitevin 1988 ³⁷		
		Benveniste 1991 ³³		
<i>multicentre</i>				
<i>external</i>				
<i>resumee: reduction of degranulation in all 3 studies</i>				
<i>isolated immune cells: basophils & histamine</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	St. Laudy 1991 ³⁸	St. Laudy 1993 ³⁹		St. Laudy 2001 ⁵⁰
		St. Laudy 1996 ⁴⁰		
		St. Laudy 1997 ⁴¹		
		St. Laudy 2006 ⁴²		
		St. Laudy 2008 ⁴³		
<i>multicentre</i>		St. Laudy 2004 ⁴⁴	Wiegant 2004 ⁴⁴	
		Ennis 2004 ⁴⁴		
		Mannaioni 2004 ⁴⁴		
<i>external</i>		Brown 2001 ⁴⁵	Guggisberg 2005 ⁴⁸	
		Lorenz 2003 ⁴⁶	Wälchli 2012 ⁴⁸	Lorenz 2003 ⁵¹
		Chirumbolo 2009 ⁴⁷		
<i>resumee: inhibition of degranulation in 13 studies, stimulation in 2, zero result in 3</i>				
<i>hypothesis: different susceptibility of basophils from different donors</i>				
<i>isolated immune cells: lymphocytes & phytolacca americana</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Colas 1984 ⁵²			
<i>multicentre</i>				
<i>external</i>			Bildet 1984 ⁵³	
<i>resumee: decrease of lymphocyte reaction in 1 study, zero result in 1</i>				
<i>isolated immune cells: lymphocytes & N-methyl-N'-nitro-N-nitrosoguanidine</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Francis 1990 ⁵⁴		Anderson 1999 ⁵⁵	
<i>multicentre</i>				
<i>external</i>				
<i>resumee: decrease of lymphocyte reaction in 1 study, zero result in 1</i>				
<i>isolated organs: rat intestine contraction & atropa belladonna or atropine sulfate</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Cristea 1997 ⁵⁶			
<i>multicentre</i>				
<i>external</i>		Schmidt 2004 ⁵⁷		
		Radau 2004 ⁵⁸		
		Michael 2004 ⁵⁹		
<i>resumee: increase or decrease of contraction at different potency levels in all 4 studies</i>				
<i>animals: amphibian metamorphosis & thyroxin / thyroindinum / triiodothyronine</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Endler 1991 ⁶⁰		Harrer 2014 ⁶⁶	
<i>multicentre</i>		Pongratz 1991 ⁶⁰	Weber 2007 ⁶²	
		van Wijk 1991 ⁶⁰		
		Zausner 2002 ^{61,68}		
		Pongratz 2002 ⁶¹		
		Lassnig 2002 ⁶¹		
		Welles 2007 ⁶²		
		Pongratz 2007 ⁶²		
		Suanjak 2007 ⁶²		
<i>external</i>		Guedes 2004 ⁶³		
		Guedes 2011 ⁶⁴		

(Continued on next page)

Table 1. (Continued)

		Harrer 2012 ⁶⁵		
<i>resumee: decrease of metamorphosis speed in 12 studies, zero result in 2</i>				
<i>hypothesis: extreme precooling of animals in Harrer 2014</i>				
<i>animals: amphibian metamorphosis & thyroxin sealed in glass vials</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Ender 1994 ^{67,69}		Dieterle 1998 ^{68,100}	
<i>multicentre</i>		Walt/Gehrer 1994 ^{67,69}		
		Pongratz 1994 ^{67,69}		
		Vinattieri 1995 ^{67,69}		
		Hilgers 1995 ^{67,69}		
		Hermann 2005 ⁶⁸		
<i>external</i>				
<i>resumee: decrease of metamorphosis speed in 6 studies, zero result in 1</i>				
<i>animals: frog climbing activity & thyroxin</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Ender 1991 ⁷⁰			
<i>multicentre</i>		Pongratz 1994 ⁷¹		
<i>external</i>				
<i>resumee: decrease of climbing activity in both studies</i>				
<i>animals: fish and homeopathic complex including iodine</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Junior 2012 ⁷²	Merlini 2014 ⁷³	Braccini 2013 ⁷⁴	
<i>multicentre</i>				
<i>external</i>				
<i>resumee: decrease in gain of size and weight in 2 studies, zero result in 1</i>				
<i>animals: arsenic trioxide poisoned mice & arsenicum album</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Mitra 1998 ⁷⁵	Mitra 1999 ⁷⁶		
		Datta 1999 ⁷⁷		
		Kundu 2000 ⁷⁸		
		Mallick 2003 ⁷⁹		
		Banerjee 2007 ⁸⁰		
		Banerjee 2008 ⁸¹		
		Banerjee 2009 ⁸²		
<i>multicentre</i>				
<i>external</i>				
<i>resumee: stimulation of damage repair in all 8 studies</i>				
<i>animals: mercury poisoned mice & mercury</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Larue 1985 ⁸³	Cal 1986 ⁸⁴		
		Larue 1986 ⁸⁵		
		Larue 1986 ⁸⁶		
<i>multicentre</i>				
<i>external</i>				
<i>resumee: protection effect in all 4 studies</i>				
<i>animals: carbon tetrachloride poisoned rats & phosphorus</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Bildet 1975 ⁸⁷			
<i>multicentre</i>				
<i>external</i>		Andresen 1985 ⁸⁸		
<i>resumee: protection effect in both studies</i>				
<i>animals: lead poisoned rats & plumbum metallicum</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Fisher 1982 ⁸⁹		Fisher 1987 ⁹⁰	
<i>multicentre</i>				
<i>external</i>				
<i>resumee: increase of excretion in 1 study, zero effect in 1</i>				
<i>animals: thrombus formation in rats & acetyl salicylic acid</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Doutremepuich 1994 ⁹¹	Belougue-Malfatti 1998 ⁹²		
		Ageujouf 2000 ⁹³		
		Eizayaga 2005 ⁹⁴		
		Doutremepuich 2007 ⁹⁵		
<i>multicentre</i>				
<i>external</i>				
<i>resumee: increase in thrombus formation in all 5 studies</i>				
<i>animals: mice under stress & gelsemium</i>				
	<i>initial study</i>	<i>repet comparable</i>	<i>repet zero</i>	<i>repet opposite</i>
<i>initial lab</i>	Bellavite 2010 ^{96,101}	Bellavite 2011 ⁹⁷		
<i>multicentre</i>				
<i>external</i>				
<i>resumee: improvement of behaviour in both studies</i>				




Figure 1 Grouping of models and studies using ultra high dilutions on enzymes, cultured cells, plants, immune cells, isolated organs, amphibians/fish and rats/mice. Columns with horizontal lines: number of models submitted to replication; columns with diagonal lines: total number of studies, i.e. including both initial and replication studies.

The selected literature base comprised 101 publications (4 of which referred to one and the same study, giving additional details)^{1–101} with data on a total of 126 studies (experiments), performed on a total of 28 experimental models. 28 of these were initial studies (on the 28 models) and 98 were replication studies. Of these replication studies, 41 were laboratory-internal replications, 28 multicentre replications and 29 external replications. Table 1 sums up the 28 models and classifies the identified 126 studies according to the type of replication and to results achieved. Multicentre studies were listed separately for the centres involved.

When all 98 replicative studies were considered, 70.4% (i.e. 69) reported a result comparable to that of the initial study, 20.4% (20) zero effect and 9.2% (9) an opposite result.

Figure 2 illustrates for the studies until 1994 (left) and the studies 1995–2015 (right) that the null-hypothesis, i.e. that there are no differences between the test and the control groups (dominance of zero results), should be rejected. Furthermore, the odds of finding a comparable result (white bars) are generally higher than of finding an opposite result (black bars). Although this is true for all three types of replication studies, it can be seen that the fraction of comparable studies diminishes from laboratory-internal (total 82.9%) to multicentre (total 75%) to external (total 48.3%).

When, for the total of the studies, the results on models that were only externally replicated (see Figure 3, left) and on models that were (mostly previous to the external replication) submitted to laboratory-internal or multicentre replication (Figure 3, right) were compared, it became obvious that the probability of an external replication producing opposite results is bigger for models that have not been further scrutinized by the initial researchers and that the probability of it producing a comparable result is bigger for models that have been submitted to laboratory-internal or multicentre replication.

Discussion

Some general issues regarding methodology and the results of a first survey as well as desirable publication standards have been discussed previously.^{102,114,115} Then, the




Figure 2 Initial and replicated studies from Table 1. Left: studies until 1994; right: 1995–2015. Ordinate: 1: initial studies; 2: internal replications; 3: multicentre; 4: external replications. Abscissa: number of studies; -: opposite result. White bars 1 are identical with the number of initial studies; white bars 2–4: comparable results; grey bars 2–4: zero effect; black bars 2–4: opposite results.




Figure 3 Initial and replicated studies from Table 1. Left: studies that were only externally replicated; right: models that were submitted to laboratory-internal or multicentre replication. For further explanation, see Figure 2.




Figure 4 Initial and replicated studies for models that have been externally reproduced with comparable results. For further explanation, see Figure 2.

point of departure in attempting an assessment of the results was one of the paradigms of homeopathy, namely that the primary aim of a homeopathic intervention is to dislodge the organism being treated from its stuck, pathological state, regardless of the direction the change may take. This paradigm finds support in clinical reports according to which healing often first manifests itself as an initial aggravation. It also finds support in zoological experiments where it seemed to be possible to steer decrease or increase of metamorphosis speed in amphibians by alternating the time interval between administration of a potentized test substance's dose.^{99,pp54–58} Indeterminacy of effect direction has been found in certain plant studies with their mutually opposite, yet in either case homogeneous, statistically significant results.^{22,24,116} The following two working hypotheses were initially considered but then found to be unsuitable for evaluating the present bibliometric data: that studies in favour of homeopathy would lead to *comparable* or *opposite* results (tested *versus* zero results), because this could lead to false-positive conclusions; or that studies in favour of a sceptic viewpoint would lead to *opposite* or *zero* results (tested *versus* comparable results), as this could lead to false-negative conclusions.

Instead, the null-hypothesis adopted was that test and control groups would not be distinguishable, i.e. there would be no result of treatment with the potency. Based on a graphic representation of the results it would then be possible to compare the odds for the occurrence of comparable results and opposite results.

Irreproducibility of results can be due to the fact that the results of the initial studies were artefacts (i.e. false-positive results). Artefacts can be due to contamination, systematic drifts or stochastic noise of the experimental set-up, which are wrongly interpreted as treatment results. Furthermore, the same reasons that can lead to result inversions may also lead to zero effects: uncontrolled relevant parameters, inappropriate outcome measures, or system inherent irreproducibility. A detailed discussion of these possible reasons for problems with reproducibility can be found elsewhere.¹¹⁷ In some cases the laboratory know-how may not have been communicated in sufficient detail. This can be obviated by organising a training phase in the

initial laboratory before an attempt is made to repeat a study. On going through the list of researchers whose replication research has at times produced zero effects one will incidentally also come across ones who can absolutely be counted among the homeopathy research community (Table 1). Another example is one external researcher who first found a comparable effect⁶⁵ and then, after joining the initial laboratory's team and, after changing a crucial laboratory parameter for exploratory purposes (i.e. temperature), a zero effect.⁶⁶

External repetitions

External replication studies yielded 48.3% comparable, 37.9% zero, and 13.8% opposite results.

We identified five models that have been reproduced by at least one external research team with comparable results:

1. Growth of wheat seedlings after treatment with potencies of silver nitrate,^{13–15}
2. Human basophil degranulation after treatment with potencies of histamine,^{38–51}
3. Highland amphibian metamorphosis after treatment with potencies of thyroxine or thyroidinum,^{60–66,98}
4. Contraction of rat intestine *in vitro* after treatment with potencies of atropa belladonna or atropine sulphate,^{56–59}
5. Experimental hepatitis of the rat due to poisoning with carbon tetrachloride after treatment with phosphorus.^{87,89}

Minor methodological differences between these replication studies and the initial studies preceding them have been critically discussed.¹⁰² While every effort was made to group the literature under survey into meaningful clusters, it is obvious that ours was not the only plausible choice and that one could find arguments in favour of other solutions. Recent studies on histological changes in cut-off tails of non-highland amphibians *Rana catesbeiana*^{63,64} were grouped together with whole organism studies on highland amphibians *Rana temporaria*^{60–62,65,66} mainly in view of the fact that both models deal with the effects of highly diluted thyroid hormones on a certain period in metamorphosis. On the other hand, some clusters could have been made even broader: For example, the cluster on the basophil model^{39–51} involving potentized histamine could have been extended to include studies on high dilutions of adrenaline¹¹⁸; or one could have created a cluster out of all studies dealing with the effects of high dilutions on DNA expression, which has been implied by studies on different systems in different laboratories¹¹⁹; or a cluster comprising studies on the anxiolytic effects of highly diluted *Gelsemium* not only on mice,^{96,97,102} but also other animals^{120–122} as well as neurocytes *in vitro*, suggesting decreased cell excitability.^{123,124} Furthermore, '>23x' may be too strict considering that in the homeopathic production process a mother tincture, depending on the pharmacopoeia used, may occasionally contain only 10^{-3} , 10^{-4} or 10^{-5} moles of the original substance per litre, i.e. their 23-fold dilution 1:10 will lead to 26x, 27x or 28x. This may be especially relevant

for the inclusion of some models by Bastide *et al.* who used 20x potencies.¹²⁵

The multicentre approach

When all replication (i.e. excluding the initial) studies are considered, 70.4% report a result comparable to that of the initial study, 20.4% a zero effect and 9.2% an opposite result. This relation is fairly well reflected by multicentre studies, i.e. studies that were centrally organised, but carried out by various researchers in different laboratories, namely 75% comparable, 14.3% zero and 10.7% opposite results. Thus, multicentre studies seem to be an adequate tool for investigating fundamental high potency models.

Laboratory-internal replication

On the other hand, initial researcher or working group studies show 82.9% comparable, 12.2% zero and 4.8% opposite results. This outcome may partly be attributable to methodological details not made explicit in the publications.

Assessing the potential of experimental models

When replication trials consistently yield zero effects one will sooner or later consider abandoning the model in question. On the other hand, when replication trials have yielded non-comparable results only in isolated cases it will make sense to review the entirety of studies published on that model before one decides to turn away from what might otherwise still grow into a fruitful branch of research.

When we screened our study material for links between the outcome of external replication studies and the history of laboratory-internal and multicentre replication work on the same model we found that initial studies that had not been followed up with laboratory-internal or multicentre replications were more likely to lead to zero effects on external replication than when they had. It thus appears that internal or multicentre replication studies, along with publications detailing the methods used and results obtained, are worth the effort if one wants to increase the probability of external researchers obtaining comparable results. On the other hand one must consider the possibility that a zero effect found on internal replication will discourage attempts at external replication.

Figure 4 shows the outcomes of the 40 studies covering the 5 models that were externally replicated with comparable results. It can be seen that some of these studies also produced zero effects or opposite results.

1994 and 2015

Furthermore, the state of replication two decades ago, i.e. in 1994,⁹⁹ was compared with that in 2015. Of the 28 models on which there had been replication work up until 2015, 15 had replication studies published before or in 1994, including 4 of the 5 promising models mentioned above. Of these 5 models, two underwent external replication before or in 1994 (Kolisko¹³ repeated by Scherer-Pon-

gratz¹⁴; Bildet⁸⁷ repeated by Andresen⁸⁸) and three after 1994 (St. Laudy,^{38–51} Cristea,^{56–59} Endler^{60–66}).

Conclusion

We found 28 experimental models in basic research on high homeopathic potencies which underwent replication research. *In total*, 24 models were replicated with comparable results, 12 models with zero effect, and 6 models with opposite results. Five models were *externally* reproduced with comparable results.

We strongly encourage further replications of published studies in order to learn more about the model systems used, to identify crucial parameters influencing experimental outcome, and to test repeatability of results. For this purpose the research methods, as well as the presentation of methods and results, should meet certain minimum standards, e.g. the guidelines for studies in homeopathy,^{98,99} either in the publication itself or in a readily available background website. As in other fields of science, a training phase in the initial laboratory may be recommendable before one attempts to repeat a study.

Acknowledgements

We thank Peter Matthießen, Stephan Baumgartner, Claudia Scherr, Karin Thieves and Christian Reich who co-authored the survey of 2009.¹⁰² We kindly acknowledge the generous supply of articles by the HOMBREX library of the Karl und Veronica Carstens Foundation.

Annotation

The authors will be grateful for comments and further information on relevant studies that fit the inclusion criteria of the bibliography. It may be of interest to the research community to further refine this publication from an annotated bibliography into a fully detailed review. It may also be of interest to use the bibliographic assessment standardized here in other fields of research, e.g. with regard to clinical trials on homeopathy.

References

- 1 Persson WM, Ginsberg AS. Die Einwirkung von Mikrodosen homöopathischer Arzneimittel (chemischer oder pflanzlicher) auf die Fermente Urease und Diastase. [The action of microdoses of homeopathic remedies (of chemicals and plants) on the enzymes urease and diastase]. *Dt Z Hom* 1932; **11**: 97–106.
- 2 Bluth M. In-vitro-Forschung mit homöopathischen Potenzen. Ein systematischer Review und eigene Versuche mit zellfreien Systemen. [In-vitro-research on homeopathic potencies. A systematic review and own experiments with cell-free systems]. Thesis. Berlin: Institut für Sozialmedizin, Epidemiologie und Gesundheitsökonomie; Charité-Universitätsmedizin, 2005.
- 3 Boyd WE. Biochemical and biological evidence of the activity of high potencies. *Br Hom J* 1954; **44**: 7–44.
- 4 Harisch G, Dittmann J. Untersuchungen zur Wirkung von Ubichinon Injeel und Injeel forte mit zellfreien Systemen. [Investigations of the result of Ubichinon Injeel and Injeel forte with cell-free systems]. *Biol Med* 1997; **26**: 99–104.

- 5 Harisch G, Dittmann J. Aktivitätsbestimmungen der sauren Phosphatase in Gegenwart von Ubichinon-Einzelpotenzien und Ubichinon-Potenzmischungen. [Determination of activity of acid phosphatase in presence of single potencies and potency mixtures of ubiquinone]. *Biol Med* 1999; **28**: 188–194.
- 6 Harisch G, Dittmann J. Unterschiedlicher Einfluß von cAMP-Potenzien und cAMP-Verdünnungen am Beispiel verschiedener Enzymsysteme. [Different influence of potencies and dilutions of cAMP exemplified on several enzyme systems]. *Biol Med* 1998; **27**: 55–62.
- 7 Harisch G, Dittmann J. Aktivitätsbestimmung der sauren Phosphatase in Gegenwart von cAMP-Einzelpotenzien und cAMP-Potenzmischungen. [Determination of activity of acid phosphatase in presence of single potencies and potency mixtures of cAMP]. *Biol Med* 1999; **28**: 4–8.
- 8 Sukul NC, Sukul A, Sinhababu SP. Potentized mercuric chloride and mercuric iodide enhance alpha-amylase activity *in vitro*. *Homeopathy* 2002; **91**: 217–220.
- 9 Carmine TC. Results of high potencies of tumour necrosis factor alpha on H₂O₂ production in cultured neuroblastoma cells by enhanced luminol-dependent chemiluminescence (ECL): a possible system for investigating the biological significance of homeopathic high potencies. *Br Hom J* 1997; **86**: 67–72.
- 10 Herberth G, Pison U. Homöopathische Arzneimittel in zellbiologischen Systemen. [Homeopathic remedies in cell-biological systems]. In: Albrecht H, Frühwald M (eds). *Jahrbuch der Karl und Veronika Carstens-Stiftung*. Essen: Band 5 (1998), 1999, pp 77–95.
- 11 Graviou E, Biron MA. Action d'une 15e CH de sulphate de cuivre sur la culture des *Chlorella vulgaris*. *Ann Hom Fr* 1971; **7**: 539–548.
- 12 Moss VA, Roberts JA, Simpson HKL. The result of copper sulphate on the growth of the alga *Chlorella*. *Br Hom J* 1977; **LXVI/3**: 169–176.
- 13 Kolisko L. *Physiologischer und physikalischer Nachweis der Wirksamkeit kleinster Entitäten bei sieben Metallen*. Dornach: Goetheanum Verlag, 1926.
- 14 Pongratz W, Endler PC. Reappraisal of a classical botanical experiment in ultra high dilution research. In: Endler PC, Schulte J (eds). *Ultra High Dilution: Physiology and Physics*. Dordrecht: Kluwer Academic Publishers, 1994, pp 121–128.
- 15 Pongratz W, Nogrask A, Endler PC. Highly diluted agitated silver nitrate and wheat seedling development. Result kinetics of a process of successive agitation phases. In: Schulte J, Endler PC (eds). *Fundamental Research in Ultra High Dilution and Homeopathy*. Dordrecht: Kluwer Academic Publishers, 1998, pp 155–187.
- 16 Betti L, Brizzi M, Nani D, Peruzzi M. Result of high dilutions of Arsenicum album on wheat seedlings from seed poisoned with the same substance. *Br Hom J* 1997; **86**: 86–89.
- 17 Brizzi M, Nani D, Peruzzi M, Betti L. Statistical analysis of the result of high dilutions of arsenic in a large dataset from a wheat germination model. *Br Hom J* 2000; **89**: 63–67.
- 18 Brizzi M, Lazzarato L, Nani D, Borghini F, Peruzzi M, Betti L. A biostatistical insight into the As₂O₃ high dilution results on the rate and variability of wheat seedling growth. *Res Complement Med – Forsch Komplementärmed* 2005; **12**: 277–283.
- 19 Nani D, Brizzi M, Lazzarato L, Betti L. The role of variability in evaluating ultra high dilution results: considerations based on plant model experiments. *Forsch Komplementärmed* 2007; **14**(5): 301–305.
- 20 Brizzi M, Elia V, Trebbi G, Nani D, Peruzzi M, Betti L. The efficacy of ultramolecular aqueous dilutions on a wheat germination model as a function of heat and aging-time. *Evid Based Complement Alternat Med* 2011. Article ID 696298.
- 21 Lahnstein L, Binder M, Thurneysen A, et al. Isopathic treatment results of arsenicum album 45x on wheat seedling growth – further reproduction trials. *Homeopathy* 2009; **98**: 198–207.
- 22 Binder M, Baumgartner S, Thurneysen A. The results of a 45x potency of arsenicum album on wheat seedling growth – a reproduction trial. *Res Complement Med – Forsch Komplementärmed* 2005; **12**: 284–291.
- 23 Pfleger A, Hofäcker J, Scherer-Pongratz W, Lothaller H, Reich C, Endler PC. The result of extremely diluted agitated gibberellic acid (10e-30) on wheat stalk growth – a two researcher pilot study. *Complement Ther Med* 2011; **19**: 164–169.
- 24 Endler PC, Matzer W, Reich C, et al. Seasonal variation of the result of extremely diluted agitated gibberellic acid (10e-30) on wheat stalk growth: a multi-researcher study. *ScientificWorldJournal* 2011; **11**: 1167–1678.
- 25 Hribar-Marko S, Graunke H, Scherer-Pongratz W, Lothaller H, Endler PC. Prestimulation of wheat seedlings with gibberellic acid followed by application of an agitated high dilution of the same hormone. *Int J High Dilution Res* 2013; **12**(42): 26–39.
- 26 Hartung H, Schiestl S, Matzer W, Endler PC. Wheat germination (20 hrs) and extremely diluted gibberellic acid (10e-30): explorative experiments on a fundamental homeopathy research model. *Eur J Integr Med* 2010; **2**: 224 [Abstract].
- 27 Kiefer P, Matzer W, Schiestl S, et al. Wheat germination and highly diluted agitated gibberellic acid (10⁻³⁰) – a multi researcher study. *Int J High Dilution Res* 2012; **11**(39): 45–59.
- 28 Scherr C, Simon M, Spranger J, Baumgartner S. Results of potentized substances on growth rate of the water plant *L. Composita*. *Complement Ther Med* 2009; **17**(2): 63–70.
- 29 Majewsky V, Scherr C, Arlt SP, et al. Reproducibility of results of homeopathically potentized gibberellic acid on the growth of *Lemna gibba* L. in a randomized and blinded bioassay. *Homeopathy* 2014; **103**: 113–126.
- 30 Baumgartner S, Doesburg P, Scherr C, Andersen JO. Development of a biocrystallisation assay for examining results of homeopathic preparations using cress seedlings. *Evid Based Complement Alternat Med* 2012; **1**–14.
- 31 Doesburg P, Andersen JO, Scherr C, Baumgartner S. Replication of specific results of a stannium metallicum 30x preparation in a cress seedling/biocrystallisation test system. In: *Proceedings 2nd HRI conference, Rome*; 2015; p. 32.
- 32 Davenas E, Beauvais F, Amara J, et al. Human basophil degranulation triggered by very dilute antiserum against IgE. *Nature* 1988; **333**: 816–818.
- 33 Benveniste J, Davenas E, Ducot B, Cornillet B, Poitevin B, Spira A. L'agitation de solutions hautement diluées n'indit pas d'activité biologique spécifique. [Agitating highly diluted solutions does not induce specific biological activity]. *CR l'Acad Paris* 1991; **312**: 461–466.
- 34 Ovelgönne JH, Bol AWJM, Hop WCJ, van Wijk R. Mechanical agitation of very dilute antiserum against IgE has no result on basophil staining properties. *Experientia* 1992; **48**: 504–508.
- 35 Hirst SJ, Hayes NA, Burridge J, Pearce FL, Foreman JC. Human basophil degranulation is not triggered by very dilute antiserum against human IgE. *Nature* 1993; **366**: 525–527.
- 36 Poitevin B, Aubin M, Benveniste J. Approche d'une analyse quantitative de l'effet d'apis mellifica sur la dégranulation des basophiles humains *in vitro*. *Innov Technol Biol Med* 1986; **7**(1): 64–68.
- 37 Poitevin B, Davenas E, Benveniste J. *In vitro* immunological degranulation of human basophils is modulated by lung histamine and *Apis mellifica*. *Br J Clin Pharmacol* 1988; **25**(4): 439–444.
- 38 Sainte-Laudy J, Belon P. Biological activity of ultra low doses. II. Result of ultra low doses of histamine on human basophil degranulation triggered by anti-IgE. In: Doutremepuich C (ed). *Ultra Low Doses*. London, Washington: Taylor & Francis, 1991, pp 139–144.
- 39 Sainte-Laudy J, Belon P. Inhibition of human basophil activation by high dilutions of histamine. *Agents Actions* 1993; **38**: 525–527.

- 40 Sainte-Laudy J, Belon P. Analysis of immunosuppressive activity of serial dilutions of histamine on human basophil activation by flow cytometry. *Inflamm Res* 1996; **45**: 33–34.
- 41 Sainte-Laudy J, Belon P. Application of flow cytometry to the analysis of the immunosuppressive result of histamine dilutions on human basophil activation: result of cimetidine. *Inflamm Res* 1997; **46**: 27–28.
- 42 Sainte-Laudy J, Belon P. Improvement of flow cytometric analysis of basophil activation inhibition by high histamine dilutions. A novel basophil specific marker: CD 203c. *Homeopathy* 2006; **95**(1): 3–8.
- 43 Sainte-Laudy J, Boujenaini N, Belon P. Confirmation of biological results of high dilutions. Results of submolecular concentrations of histamine and 1-, 3- and 4-methylhistamines on human basophil activation. *Inflamm Res* 2008; **57**(S1): S01–S02.
- 44 Belon P, Cumps J, Ennis M, et al. Histamine dilutions modulate basophile activation. *Inflamm Res* 2004; **53**: 181–188.
- 45 Brown V, Ennis M. Flow-cytometric analysis of basophil activation: inhibition by histamine at conventional and homeopathic concentrations. *Inflamm Res* 2001; **50**: 47–48.
- 46 Lorenz I, Schneider EM, Stolz P, Brack A, Strube J. Sensitive flow cytometric method to test basophil activation influenced by homeopathic histamine dilutions. *Forsch Komplementärmed Klass Naturheilkd* 2003; **10**: 316–324.
- 47 Chirumbolo S, Brizzi M, Ortolani R, Vella A, Bellavite P. Inhibition of CD203c membrane up-regulation in human basophils by high dilutions of histamine: a controlled replication study. *Inflamm Res* 2009; **58**: 755–764.
- 48 Guggisberg AG, Baumgartner S, Tschopp CM, Heusser P. Replication study concerning the results of homeopathic dilutions of histamine on human basophil degranulation *in vitro*. *Complement Ther Med* 2005; **13**: 91–100.
- 49 Wälchli C, Baumgartner S. Flow cytometric analysis of the inhibition of human basophil activation by histamine high dilutions – a replication study. *Int J High Dilution Res* 2012; **11**(40): 172–173.
- 50 Sainte-Laudy J. Stimulatory result of high dilutions of histamine on activation of human basophils induced by anti-IgE. *Inflamm Res* 2001; **50**: 63–64.
- 51 Lorenz I, Schneider EM, Stolz P, Brack A, Strube J. Influence of the diluent on the result of highly diluted histamine on basophil activation. *Homeopathy* 2003; **92**: 11–18.
- 52 Colas H, Aubin M, Picard P, Lebecq JC, Bastide JM. Inhibition du test de transformation lymphoblastique (TTL) à la phytohématagglutinine (PHA) par *Phytolacca americana* en dilutions homéopathiques. [Inhibition of lymphoblast transformation test (LTT) by phythemagglutinine (PHA) through *Phytolacca americana* in homeopathic dilutions]. *Hom Franc* 1984; **72**: 219–224.
- 53 Bildet J, Dupont H, Aubin M, et al. Action *in vitro* de dilutions infinitésimales de *Phytolacca americana* sur la transformation lymphoblastique (TTL) à la phytohématagglutinine. [In vitro action of infinitesimal dilutions of *Phytolacca americana* on the transformation of lymphoblasts by phytohemagglutinine]. *Hom Franc* 1984; **72**: 225–230.
- 54 Francis AJ, Anderson D, Fisher P. Further studies of the ‘adaptive’ repair response in human lymphocytes after treatment with MNNG. *Environ Moloc Mutagen* 1990; **15**(S17): 69 [Abstract].
- 55 Anderson D, Edwards AJ, Fisher P, Lovell DP. Statistical analysis of adaptive response in sister chromatid exchanges in human lymphocytes after treatment with very low and extremely low doses of N-methyl-N'-nitro-N-nitrosoguanidine using a study design to control variability. *Br Hom J* 1999; **88**(1): 7–16.
- 56 Cristea A, Nicula S, Darie V. Pharmacodynamic results of very high dilutions of belladonna on the isolated rat duodenum. In: Bastide M (ed). *Signals and Images*. Dordrecht: Kluwer Academic Publishers, 1997, pp 161–170.
- 57 Schmidt F, Süß WG, Nieber K. *In-vitro* Testung von homöopathischen Verduennungen. *Biol Med* 2004; **1**: 32–37.
- 58 Radau K. *Materialwissenschaftliche Untersuchungen an pharmazeutischen Hilfsstoffen und ihre Bedeutung für die Herstellung homöopathischer Arzneimittel*. Thesis. Fakultät für Biowissenschaften, Pharmazie und Psychologie, Universität Leipzig, 2004.
- 59 Michael S. *Untersuchungen zur Wirkung von homöopathischen Verdünnungen am isolierten Ileum der Ratte*. Thesis. Institut für Pharmazie, Universität Leipzig, 2004.
- 60 Ender PC, Pongratz W, van Wijk R, Kastberger G, Haidvogel M. Results of highly diluted succussed thyroxine on metamorphosis of highland frogs. *Berl J Res Hom* 1991; **1**: 151–160.
- 61 Zausner C, Lassnig H, Ender PC, et al. Die Wirkung von homöopathisch zubereitetem Thyroxin auf die Metamorphose von Hochlandamphibien. Ergebnisse einer multizentrischen Kontrollstudie. *Perfusion* 2002; **17**: 268–276.
- 62 Welles SU, Suanjak-Traidl E, Weber S, et al. Pretreatment with thyroxine (10e-8) and the result of homeopathically prepared thyroxine (10e-30) on highland frogs – a multi-researcher study. *Res Complement Med – Forsch Komplementärmed* 2007; **14**: 353–357.
- 63 Guedes JRP, Ferreira CM, Guimaraes HMB, Saldiva PHN, Capelozzi VL. Homeopathically prepared dilution of *Rana catesbeiana* thyroid glands modifies its rate of metamorphosis. *Homeopathy* 2004; **93**: 132–137.
- 64 Guedes JR, Carrasco S, Ferreira CM, et al. Ultra high dilution of triiodothyronine modifies cellular apoptosis in *Rana catesbeiana* tadpole tail *in vitro*. *Homeopathy* 2011 Oct; **100**(4): 220–227.
- 65 Harrer B. Replication of an experiment on extremely diluted thyroxine and highland amphibians. *Homeopathy* 2013 Jan; **102**(1): 25–30 <http://dx.doi.org/10.1016/j.homp.2012.09.003>.
- 66 Harrer B, Ender PC. *Highland amphibians and extremely diluted thyroxine – pre-experiment storage at low temperature (4°C)*. HAR/A, 1–2, www.ResProceedingsIUC.net; 2013.
- 67 Ender PC, Pongratz W, Smith CW, Schulte J. Non-molecular information transfer from thyroxine to frogs with regard to ‘homoeopathic’ toxicology. *Vet Hum Toxicol* 1995; **37**(3): 259–260.
- 68 Hermann B. *Zur Wirkung von ‘homöopathisch’ zubereitetem Thyroxin (10e-30) in Glasphiolen auf die Metamorphose vorstimulierter Rana temporaria-Larven*. Graz: Interuniversitäres Kolleg, 2005.
- 69 Ender PC, Heckmann C, Lauppert E, et al. The metamorphosis of amphibian and information of thyroxine storage via the bipolar fluid water. In: Schulte J, Ender PC (eds). *Fundamental Research in Ultra High Dilution and Homeopathy*. Dordrecht: Kluwer Academic Publishers, 1998.
- 70 Ender PC, Pongratz W, Kastberger G, Wiegant FAC, Haidvogel M. Climbing activity in frogs and the result of highly diluted succussed thyroxine. *Br Hom J* 1991; **80**: 194–200.
- 71 Ender PC, Pongratz W, Kastberger G, Wiegant FAC, Schulte J. The result of highly diluted agitated thyroxine on the climbing activity of frogs. *J Vet Hum Toxicol* 1994; **36**: 56–59.
- 72 Junior RP, Vargas L, Valentim-Zabott M, Ribeiro RP, da Silva AV, Otutumi LK. Morphometry of white muscle fibers and performance of Nile tilapia (*Oreochromis niloticus*) fingerlings treated with methyltestosterone or a homeopathic complex. *Homeopathy* 2012; **101**: 154–158.
- 73 Merlini LS, Vargas L, Piau R, Ribeiro RP, Merlini NB. Results of a homeopathic complex on the performance and cortisol levels in Nile tilapia (*Oreochromis niloticus*). *Homeopathy* 2014; **103**: 139–142.
- 74 Braccini GL, Natali MR, Ribeiro RP, et al. Morpho-functional response of Nile tilapia (*Oreochromis niloticus*) to a homeopathic complex. *Homeopathy* 2013; **102**: 233–241.
- 75 Mitra K, Kundu SN, Khuda-Bukhsh AR. Efficacy of a potentized homeopathic drug (Arsenicum Album-30) in reducing toxic results produced by arsenic trioxide in mice: I. On rate of accumulation of arsenic in certain organs. *Complement Ther Med* 1998; **6**: 178–184.

- 76 Mitra K, Kundu SN, Khuda-Bukhsh AR. Efficacy of a potentized homeopathic drug (Arsenicum Album-30) in reducing toxic results produced by arsenic trioxide in mice: II. On alterations of body weight, tissue weight and total protein. *Complement Ther Med* 1999; **7**: 24–34.
- 77 Datta S, Mallick P, Bukhsh AR. Efficacy of a potentized homeopathic drug (Arsenicum Album-30) in reducing genotoxic results produced by arsenic trioxide in mice. *Complement Ther Med* 1999; **7**(2): 62–75. and **7**(3): 156–163.
- 78 Kundu SN, Mitra K, Khuda Bukhsh AR. Efficacy of a potentized homeopathic drug (Arsenicum-album-30) in reducing cytotoxic results produced by arsenic trioxide in mice. *Complement Ther Med* 2000; **8**(2): 76–81. and **8** (3): 157–165.
- 79 Mallick P, Chakrabarti Mallick J, Guha B, Khuda-Bukhsh AR. Ameliorating result of microdoses of a potentized homeopathic drug, arsenicum album, on arsenic-induced toxicity in mice. *BMC Complement Alternat Med* 2003; **3**: 7.
- 80 Banerjee P, Biswas SJ, Belon P, Khuda-Bukhsh AR. A potentized homeopathic drug, Arsenicum Album 200, can ameliorate genotoxicity induced by repeated injections of arsenic trioxide in mice. *J Vet Med A Physiol Pathol Clin Med* Sep 2007; **54**(7): 370–376.
- 81 Banerjee P, Bhattacharyya SS, Pathak S, Naoual B, Belon P, Khuda-Bukhsh AR. Comparative efficacy of two microdoses of a potentized homeopathic drug, arsenicum album, to ameliorate toxicity included by repeated sublethal injections of arsenic trioxide in mice. *Pathobiology* 2008; **75**: 156–170.
- 82 Banerjee P, Bhattacharyya SS, Pathak S, Boujedaini N, Belon P, Khuda-Bukhsh AR. Evidences of protective potentials of microdoses of ultra-high diluted arsenic trioxide in mice receiving repeated injections of arsenic trioxide. *Evid Based Complement Alternat Med* 2009 [in press].
- 83 Larue F, Cal JC, Tetau M, Buisard AM, Guillemain J, Cambar J. Mise en évidence de l'effet Protecteur de différentes dilutions de mercurius corrosivus. *Cah Biothér* 1985; **88**: 71–74.
- 84 Cal JC, Larue F, Guillemain J, Cambar J. Chronobiological approach of protective result of mercurius corrosivus against mercury-induced nephrotoxicity. *Annu Rev Chronopharmacol* 1986; **3**: 99–103.
- 85 Larue F, Cal JC, Guillemain J, Cambar J. Influence de la durée de prétraitement sur l'effet de mercurius corrosivus vis-à-vis de la toxicité induite par le chlorure mercurique chez la souris. *Homéopathie Française* 1986; **74**(5): 275–281.
- 86 Larue F, Cal JC, Guillemain J, Cambar J. Variations saisonnières et circadiennes de l'efficacité du prétraitement par mercurius corrosivus 15CH vis-à-vis de la toxicité induite par le chlorure mercurique. *Bull du Groupe d'Etude des Rhythm Biol* 1986; **18**(1–2): 8–9.
- 87 Bildet J. *Etude de l'action de différentes dilutions homéopathiques de phosphore blanc (Phosphorus) sur l'hépatite toxique du rat*. Thèse. Université de Bordeaux II, 1975.
- 88 Andresen M. *Zytosolische und mitochondriale Effekte einer Intoxikation mit CCl4 am Beispiel des Lebergewebes der Ratte. Einfluss von Phosphorus D6 und Phosphorus D30*. Thesis. Tierärztliche Hochschule Hannover, 1985.
- 89 Fisher P. The treatment of experimental lead intoxication in rats by penicillinamine and plumbum met. In: *Proceedings of the 35th Congress of the Liga Medicorum Homeopathica Internationalis, Brighton*; 1982; pp 320–332.
- 90 Fisher P, House I, Belon P, Turner P. The influence of the homeopathic remedy Plumbum metallicum on the excretion kinetics of lead in rats. *Hum Toxicol* 1987; **6**: 321–324.
- 91 Doutremepuich C, Aguejouf O, Pintigny D, Sertillanges MN, De Seze O. Thrombogenic properties of ultra-low-dose of acetylsalicylic acid in a vessel model of laser-induced thrombus formation. *Thromb Res* 1994; **76**: 225–229.
- 92 Belougne-Malfatti E, Aguejouf O, Doutremepuich F, Belon P, Doutremepuich C. Combination of two doses of acetyl salicylic acid: experimental study of arterial thrombosis. *Thromb Res* 1998; **90**: 215–221.
- 93 Aguejouf O, Malfatti E, Belon P, Doutremepuich C. Time related neutralization of two doses acetyl salicylic acid. *Thromb Res* 2000; **100**(4): 317–323.
- 94 Eizayaga FX, Aguejouf O, Belon P, Doutremepuich C. Platelet aggregation in portal hypertension and its modification by ultra-low doses of aspirin. *Pathophysiol Haemost Thromb* 2005; **34**(1): 29–34.
- 95 Doutremepuich C, Aguejouf O, Eizayaga FX, Desplat V. Reverse result of aspirin: is the prothrombotic result after aspirin discontinuation mediated by cyclooxygenase 2 inhibition? *Pathophysiol Haemost Thromb* 2007; **36**(1): 40–44.
- 96 Magnani P, Conforti A, Zanolin E, Marzotto M, Bellavite P. Dose-result study of *Gelsemium sempervirens* in high dilutions on anxiety-related responses in mice. *Psychopharmacol Berl* 2010 Jul; **210**(4): 533–545.
- 97 Bellavite P, Magnani P, Zanolin E, Conforti A. Homeopathic doses of *Gelsemium sempervirens* improve the behavior of mice in response to novel environments. *Evid Based Complement Alternat Med* 2011; **2011**: 362517.
- 98 Zausner-Lukitsch C. *Auswirkungen von homöopathisch zubereitetem Thyroxin auf die Metamorphosegeschwindigkeit von Rana temporaria unter besonderer Berücksichtigung der Einzelhaltung und unterschiedlicher Methoden der Applikation*. Thesis. Vienna University, 2001.
- 99 Endler PC, Schulte J (eds). *Ultra High Dilution: Physiology and Physics*. Dordrecht: Kluwer Academic Publishers, 1994.
- 100 Dieterle D. *Überprüfung einer Hypothese zum indirekten Einfluß potenziierter Thyroxinlösungen auf die Metamorphosegeschwindigkeit von Rana temporaria*. Thesis. University of Tübingen, 1999.
- 101 Bellavite P, Conforti A, Marzotto M, et al. Testing homeopathy in mouse emotional response models: pooled data analysis of two series of studies. *Evid Based Complement Alternat Med* 2012; **2012**: 954374.
- 102 Endler PC, Thieves K, Reich C, et al. Replications of fundamental research models for homeopathically prepared dilutions beyond 10e-23: a bibliometric study. *Homeopathy* 2010; **99**: 25–36.
- 103 Forschung im Dienste der Gesundheit (Projektträgerschaft). *Unkonventionelle Medizinische Richtungen. Bestandsaufnahme zur Forschungssituation*. Bonn: Wirtschaftsverlag, Verlag für neue Wissenschaft, 1992.
- 104 Linde K, Jonas WB, Melchart D, Worku F, Wagner H, Eitel F. Critical review and meta-analysis of serial agitated dilutions in experimental toxicology. *Hum Exp Toxicol* 1994; **13**: 481–482.
- 105 Endler PC. *Homeopathy research – an expedition report*. Edition Interuniversity College, Graz 2003/Expedition Homöopathieforschung – Ein altes Heilsystem wird plausibel. Vienna: Maudrich Verlag, 1998.
- 106 Endler PC. Fundamental research into high dilution results. A classification of non-clinical research topics. In: Schulte J, Endler PC (eds). *Fundamental Research in Ultra High Dilution and Homeopathy*. Dordrecht: Kluwer Academic Publishers, 1998.
- 107 Vickers AJ. External replication of pre-clinical research in homeopathy: a systematic review. *Res Complement Med – Forsch Komplementärmed* 1999; **6**: 311–320.
- 108 Albrecht H, van Wijk R, Dittloff S. A new database on basic research in homeopathy. *Homeopathy* 2002; **91**: 162–165.
- 109 Baumgartner S. Reproductions and reproducibility in homeopathy: dogma or tool? *J Altern Complement Med* 2005; **11**(5): 771–772.
- 110 Walach H, Jones WB, Ives J, Wijk RV, Weingartner O. Research on homeopathy: state of the art. *J Altern Complement Med* 2005; **11**: 813–829.
- 111 Göldner C. *Review und Evaluierung von neuen, mit homöopathischen Zubereitungen durchgeführten toxikologischen Experimenten*. Thesis. University of Graz, 2006.

- 112 Witt CM, Bluth M, Albrecht H, Weissshuhn TE, Baumgartner S, Willich SN. The *in vitro* evidence for an result of high homeopathic potencies-a system review of the literature. *Complement Ther Med* 2007; **15**: 128–138.
- 113 Baumgartner S. The state of basic research on homeopathy. In: Albrecht H, Witt C (eds). *New Directions in Homeopathy Research – Advice from an Interdisciplinary Conference*. Essen: KVC-Verlag, 2009, pp 107–130.
- 114 Stock-Schröer B, Albrecht H, Betti L, et al. Reporting experiments in homeopathic basic research (REHBaR) – description of the checklist development. *eCAM* 2009; <http://dx.doi.org/10.1093/ecam/nep170>.
- 115 Stock-Schröer B, Albrecht H, Betti L, et al. Reporting experiments in homeopathic basic research (REHBaR) – a guideline for authors. *Homeopathy* 2009; **98**: 287–298.
- 116 Bell IR, Baldwin CM, Schwartz GE. Translating a nonlinear systems theory model for homeopathy into empirical tests. *Altern Ther Health Med* 2002; **8**(3): 58–66.
- 117 Bellavite P. Complexity science and homeopathy. A synthetic overview. *Homeopathy* 2003; **92**: 203–212.
- 118 Mannaioni PF, Mastroianni R, Mastrangelo D. Adrenaline inhibits the immunological activation of human basophils at pharmacological and ultra-low doses. *Med Sci Monit* 2010; **16**: BR227–BR232.
- 119 Bellavite P, Signorini A, Marzotto M, Moratti E, Bonafini C, Olioso D. Cell sensitivity, non-linearity and inverse effects. *Homeopathy* 2015; **104**: 139–160.
- 120 Bellavite P, Magnani P, Marzotto M, Conforti A. Assays of homeopathic remedies in rodent behavioural and psychopathological models. *Homeopathy* 2009; **98**: 208–227.
- 121 Venard C, Boujedaini N, Mensah-Nyagan AG, Patte-Mensah C. Comparative analysis of gelsemine and *Gelsemium sempervirens* activity on neurosteroid allopregnanolone formation in the spinal cord and limbic system. *Evid Based Complement Alternat Med* 2011; **2011**: 407617.
- 122 Meyer L, Boujedaini N, Patte-Mensah C, Mensah-Nyagan AG. Pharmacological effect of gelsemine on anxiety-like behavior in rat. *Behav Brain Res* 2013; **253**: 90–94.
- 123 Olioso D, Marzotto M, Moratti E, Brizzi M, Bellavite P. Effects of *Gelsemium sempervirens* L. on pathway-focused gene expression profiling in neuronal cells. *J Ethnopharmacol* 2014; **153**: 535–539.
- 124 Marzotto M, Olioso D, Brizzi M, Tononi P, Cristofolletti M, Bellavite P. Extreme sensitivity of gene expression in human SH-SY5Y neurocytes to ultra-low doses of *Gelsemium sempervirens*. *BMC Complement Alternat Med* 2014; **14**: 104.
- 125 Bonamin LV, Bellavite P. Immunological models in high dilution research following M Bastide. *Homeopathy* 2015; **104**: 263–268.